

Fermions tunneling from higher-dimensional Reissner-Nordström black hole

Kai Lin

E-mail: lk314159@126.com

ShuZheng Yang

E-mail: szyangcwnu@126.com

Institute of Theoretical Physics, China West Normal University, NanChong, SiChuan 637002 China

Abstract.

Based on semiclassical tunneling method, our focus is charged fermions tunneling from higher-dimensional Reissner-Nordström black hole. We first simplify the Dirac equation via semiclassical approximation, and then obtain a semiclassical Hamilton-Jacobi equation. Using the Hamilton-Jacobi equation, we study the Hawking temperature and fermions tunneling rate at the event horizon of the higher-dimensional Reissner-Nordström black hole.

Keywords: Higher Dimensional Reissner-Nordström Black Hole, Hawking radiation, Dirac Equation

PACS numbers: 04.70.Dy, 04.62.+v, 03.65.Sq

Hawking radiation is an important prediction in modern gravitation theory [1–5]. Recently, Kraus, Parikh and Wilczek proposed quantum tunneling theory to explain and study Hawking radiation [6–31], and then Semiclassical Hamilton-Jacobi method is put forward to research the properties of scalar particles' tunnels [32–36]. In 2007, Kerner and Mann investigated the 1/2 spin fermion tunneling from static black holes [37]. In their work, the spin up and spin down cases are researched respectively, and the radial equations are obtained, so that they can finally determine the Hawking temperature and tunneling rate at the event horizon. Subsequently, Kerr and Kerr-Newman black holes cases, the charged dilatonic black hole case, the de Sitter horizon case, the BTZ black hole case, 5-dimensional spacetime cases and several non-stationary black hole cases were all researched respectively [38–48], and we used Hamilton-Jacobi method to study the fermion tunneling from higher-dimensional uncharged black holes [49–50]. However, up to now, no one has studied higher-dimensional charged black holes cases, so we set out to research that case. In our work, we developed the Kerner and Mann method, and proved the semiclassical Hamilton-Jacobi equation not only can be obtained with the Klein-Gordon equation of curved spacetime, but also with the Dirac equation in curved spacetime. Applying the Hamilton-Jacobi equation, we can then obtain semiclassical Hawking temperature and tunneling rate at the event horizon of higher-dimensional Reissner-Nordström black hole.

In modern physics theory, the concept of an extra dimension can help to solve some theoretical issues, so several higher dimensional metrics of curved spacetime was put forward. The metric of static charged (n+2)-dimensional Reissner-Nordström black hole is given by [10,51–55]

$$ds^2 = -f(r)dt^2 + f^{-1}(r)dr^2 + r^2 d\Omega_n^2 \quad (1)$$

where $d\Omega_n^2$ is the metric of n-dimensional sphere

$$d\Omega_n^2 = \sum_{i=1}^n h^{ii} d\theta_i^2 = d\theta_1^2 + \sin^2\theta_1 d\theta_2^2 + \sin^2\theta_1 \sin^2\theta_2 d\theta_3^2 + \dots + \prod_{i=1}^{n-1} \sin^2\theta_i d\theta_n^2 \quad (2)$$

and

$$f(r) = 1 - \frac{\omega_n M}{r^{n-1}} + \frac{\omega_n Q^2}{2(n-1)V_n r^{2n-2}} \quad \omega_n = \frac{16\pi}{nV_n} \quad (3)$$

M and Q are mass and electric charge of black hole, and the electro-magnetic potential is

$$A_\mu = \left(\frac{Q}{(n-1)V_n r^{n-1}}, 0, 0, 0 \right) \quad (4)$$

where V_n is volume of unit n-sphere (we can adopt the units $G = c = \hbar = 1$). The outer/ inner horizon located at

$$r_{\pm}^{n-2} = \frac{\omega_n}{2} \left[M \pm \sqrt{M^2 - \frac{nQ^2}{8\pi(n-1)}} \right] \quad (5)$$

Obviously, at the horizons, the equation $f(r_{\pm}) = 0$ should be satisfied. However, the physical property near the inner horizon cannot be researched, so we just study the fermion tunneling

at the outer event horizon of this black hole. The charged Dirac equation in curved spacetime is

$$\gamma^\mu D_\mu \Psi + \frac{m}{\hbar} = 0 \quad \mu = 1, 2, 3, \dots, n+2 \quad (6)$$

where

$$D_\mu = \partial_\mu + \frac{i}{2} \Gamma^\alpha{}_\mu{}^\beta + \frac{iqA_\mu}{\hbar} \quad (7)$$

$$\Pi_{\alpha\beta} = \frac{i}{4} [\gamma_\alpha, \gamma_\beta] \quad (8)$$

m and q are mass and electric charge of the particles. The gamma matrices in curved spacetime need to be satisfied

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} I \quad (9)$$

After the gamma matrices are defined, we choose the gamma matrices in $(n+2)$ -dimensional flat spacetime as

$$\tilde{\gamma}_{m \times m}^1 = \begin{pmatrix} I_{\frac{m}{2} \times \frac{m}{2}} & 0 \\ 0 & -I_{\frac{m}{2} \times \frac{m}{2}} \end{pmatrix} \quad (10)$$

$$\tilde{\gamma}_{m \times m}^2 = \begin{pmatrix} 0 & I_{\frac{m}{2} \times \frac{m}{2}} \\ I_{\frac{m}{2} \times \frac{m}{2}} & 0 \end{pmatrix} \quad (11)$$

$$\tilde{\gamma}_{m \times m}^\eta = \begin{pmatrix} 0 & i\tilde{\gamma}_{\frac{m}{2} \times \frac{m}{2}}^{\eta-2} \\ -i\tilde{\gamma}_{\frac{m}{2} \times \frac{m}{2}}^{\eta-2} & 0 \end{pmatrix} \quad \eta = 3, 4, 5, \dots, n+2 \quad (12)$$

where $I_{\frac{m}{2} \times \frac{m}{2}}$ and $\tilde{\gamma}_{\frac{m}{2} \times \frac{m}{2}}^\eta$ are unit matrices and flat gamma matrices with $\frac{m}{2} \times \frac{m}{2}$ order, and $m = 2^{(n+2)/2}$ ($m = 2^{(n+1)/2}$) is the order of the matrices in even (odd) dimensional space-time. Corresponding to the flat case, the gamma matrices can be chosen as

$$\gamma_{m \times m}^t = \frac{i}{\sqrt{f}} \tilde{\gamma}_{m \times m}^1 \quad (13)$$

$$\gamma_{m \times m}^r = \sqrt{f} \tilde{\gamma}_{m \times m}^2 \quad (14)$$

$$\gamma_{m \times m}^\eta = r^{-1} \sqrt{h^{\eta\eta}} \tilde{\gamma}_{m \times m}^\eta \quad \eta = 3, 4, 5, \dots, n+2. \quad (15)$$

Now, let's simplify the Dirac equation via semiclassical approximation, and rewrite the spinor function as

$$\Psi = \begin{pmatrix} A_{\frac{m}{2} \times 1}(t, r, \dots, x^\eta, \dots) \\ B_{\frac{m}{2} \times 1}(t, r, \dots, x^\eta, \dots) \end{pmatrix} e^{\frac{i}{\hbar} S(t, r, \dots, x^\eta, \dots)} \quad (16)$$

where $A_{\frac{m}{2} \times 1}(t, r, \dots, x^\eta, \dots)$ and $B_{\frac{m}{2} \times 1}(t, r, \dots, x^\eta, \dots)$ are column matrices with $\frac{m}{2} \times 1$ order, and S is classical action. Via semiclassical approximation method, Substituting Eq.(16) into Eq.(6) and dividing the exponential term and multiplying by $\hbar$, we can get

$$\begin{pmatrix} C & D \\ E & F \end{pmatrix} \begin{pmatrix} A_{\frac{m}{2} \times 1} \\ B_{\frac{m}{2} \times 1} \end{pmatrix} = 0 \quad (17)$$

$$C = -\frac{1}{\sqrt{f}} \left(\frac{\partial S}{\partial t} + qA_t \right) I_{\frac{m}{2} \times \frac{m}{2}} + m I_{\frac{m}{2} \times \frac{m}{2}} \quad (18)$$

$$D = i \sqrt{f} \frac{\partial S}{\partial r} I_{\frac{m}{2} \times \frac{m}{2}} - \sum_{\eta} r^{-1} \sqrt{h^{\eta\eta}} \frac{\partial S}{\partial x^{\eta}} \tilde{\gamma}^{\eta-2}_{\frac{m}{2} \times \frac{m}{2}} \quad (19)$$

$$E = i \sqrt{f} \frac{\partial S}{\partial r} I_{\frac{m}{2} \times \frac{m}{2}} + \sum_{\eta} r^{-1} \sqrt{h^{\eta\eta}} \frac{\partial S}{\partial x^{\eta}} \tilde{\gamma}^{\eta-2}_{\frac{m}{2} \times \frac{m}{2}} \quad (20)$$

$$F = \frac{1}{\sqrt{f}} \left(\frac{\partial S}{\partial t} + qA_t \right) I_{\frac{m}{2} \times \frac{m}{2}} + m I_{\frac{m}{2} \times \frac{m}{2}} \quad (21)$$

Solving Eq.(17), we have

$$(E - FD^{-1}C)A_{\frac{m}{2} \times 1} = 0 \quad (22)$$

$$(F - EC^{-1}D)B_{\frac{m}{2} \times 1} = 0 \quad (23)$$

It is evident that the coefficient matrices of (22) and (23) must vanish, when $A_{\frac{m}{2} \times 1}$ and $B_{\frac{m}{2} \times 1}$ have non-trivial solutions. Due to the fact that $CD = DC$, we can write the condition that determinant of coefficient vanish as

$$\det(ED - FC) = 0 \quad (24)$$

From the relation of flat gamma matrices $\{\tilde{\gamma}^{\mu}, \tilde{\gamma}^{\nu}\} = 2\delta_{\mu\nu}$, we can obtain the semiclassical Hamilton-Jacobi equation in $(n+2)$ -dimensional Reissner-Nordström space time

$$-\frac{1}{f} \left(\frac{\partial S}{\partial t} + qA_t \right)^2 + f \left(\frac{\partial S}{\partial r} \right)^2 + \dots + g^{\eta\eta} \left(\frac{\partial S}{\partial x^{\eta}} \right)^2 + \dots + m^2 = 0 \quad (25)$$

Using the Hamilton-Jacobi equation, in charged static spacetime, we can separate the variables for the action as

$$S = -\omega t + R(r) + Y(\dots, x^{\eta}, \dots) + K \quad (K \text{ is a constant}) \quad (26)$$

and the Hamilton-Jacobi equation is broken up as

$$-\frac{1}{f} (\omega - qA_t)^2 + f \left(\frac{dR}{dr} \right)^2 + m^2 = \frac{\lambda}{r^2} \quad (27)$$

$$\sum_{\eta} h^{\eta\eta} \left(\frac{\partial Y}{\partial x^{\eta}} \right)^2 + \lambda = 0 \quad (28)$$

where Eq.(27) and Eq.(28) are radial and non-radial equations respectively, and λ is a constant. However, we only research on the radial equation, because the tunnel at the event horizon is radial. From Eq.(27), we can get

$$\frac{dR(r)}{dr} = \pm \frac{\sqrt{(\omega - qA_t)^2 r^2 + f(\lambda - m^2 r^2)}}{fr} \quad (29)$$

Near the event horizon, the radial action is given by

$$ImR_{\pm} = \pm \frac{\pi(\omega - \omega_0)}{f'(r_+)} + ImK \quad (30)$$

where, R_+ is part of the outgoing solution, while R_- is the part of incoming solution, and

$$\omega_0 = q \frac{Q}{(n-1)V_n r_+^{n-1}} \quad (31)$$

So the tunneling rate is

$$\Gamma = \frac{\text{Prob[out]}}{\text{Prob[in]}} = \frac{\exp(-2\text{Im}S_+)}{\exp(-2\text{Im}S_-)} = \frac{\exp(-2\text{Im}R_+ + \text{Im}K)}{\exp(-2\text{Im}R_- + \text{Im}K)} = \exp\left(\frac{-4\pi(\omega - \omega_0)}{f'(r_+)}\right) \quad (32)$$

where Im represents the imaginary part of the function, and the Hawking temperature is

$$T_H = \frac{f'(r_+)}{4\pi} \quad (33)$$

Of course, In this paper, we researched fermions tunneling from higher-dimensional Reissner-Nordström black holes. We got the Hamilton-Jacobi equation, and this work shows that semiclassical Hamilton-Jacobi equation can describe the property of both 0 spin scalar particles and 1/2 spin fermions. In this work, we did not emphasize dimensions of spacetime larger than (3+1) dimensions, so the method should be used in the research of (3+1)-dimensions and lower cases. However, above work base on the semiclassical approximation. Recently, Banerjee and Majhi proposed a new method beyond semiclassical approximation to research the quantum tunneling, and their work show that the conclusion should be corrected [56–65]. Our further work will generalize the semiclassical Hamilton-Jacobi method into the method beyond semiclassical approximation.

Acknowledgements: This work is supported by National Natural Science Foundation of China (No.10773008)

References

- [1] S. W. Hawking, *Nature* **248** 30 (1974)
- [2] S. W. Hawking, *Commun.Math.Phys.* **43** 199 (1975)
- [3] S. P. Robinson and F. Wilczek, *Phys.Rev.Lett.* **95** 011303 (2005) [arXiv:gr-qc/0502074]
- [4] T. Damour and R. Ruffini, *Phys.Rev.D* **14** 332 (1976)
- [5] S. Sannan, *Gen.Relativ.Gravit.* **20** 239 (1988)
- [6] P. Kraus and F. Wilczek 1995 *Nucl.Phys.B* **433** 403 [arXiv:gr-qc/9408003]
- [7] M. K. Parikh and F. Wilczek, *Phys.Rev.Lett.* **85** 5042 (2000) [arXiv:hep-th/9907001]
- [8] S. Hemming and E. Keski-Vakkuri, *Phys. Rev. D* **64** 044006 (2001)[arXiv:gr-qc/0005115]
- [9] Q. Q. Jiang, S. Q. Wu and X. Cai, *Phys.Rev.D* **75**, 064029 (2007)
- [10] S. Q. Wu and Q. Q. Jiang [arXiv:hep-th/0603082]
- [11] R. Kerner and R. B. Mann, *Phys.Rev.D* **73** 104010 (2006)
- [12] J. Ren, J. Y. Zhang and Z Zhao, *Chin.Phys.Lett.* **23** 2019 (2006)
- [13] S. Iso, H. Umetsu, and F. Wilczek, *Phys.Rev.D* **74** 044017 (2006)[hep-th/0606018]
- [14] S. Iso, T. Morita, and H. Umetsu, *J.High.Energy.Phys.* **0704** 068 (2007) [hep-th/0612286]
- [15] S. Z. Yang, H. L. Li, Q. Q. Jiang and M. Q. Liu, *Sic.In.Chin.G* **50** 249 (2007)
- [16] E. C. Vagenas, *Phys.Lett.B* **503** 399 (2001) [arXiv:hep-th/0012134]
- [17] E. C. Vagenas, *Mod.Phys.Lett.A* **17** 609 (2002) [arXiv:hep-th/0108147]
- [18] E. C. Vagenas, *Phys.Lett.B* **533** 302 (2002) [arXiv:hep-th/0109108]
- [19] A. J. M. Medved, *Class.Quant.Grav.* **19** 589 (2002) [arXiv:hep-th/0110289]
- [20] M. K. Parikh, *Phys.Lett.B* **546** 189 (2002) [arXiv:hep-th/0204107]
- [21] A. J. M. Medved, *Phys.Rev.D* **66** 124009 (2002) [arXiv:hep-th/0207247]

- [22] E. C. Vagenas, *Phys.Lett.B* **559** 65 (2003) [arXiv:hep-th/0209185]
- [23] M. K. Parikh, [arXiv:hep-th/0402166]
- [24] J. Y. Zhang and Z. Zhao 2006 *Acta.Phys.Sin.* **55** 3796 (in Chinese)
- [25] J. Y. Zhang and Z. Zhao 2006 *Phys.Lett.B* **638** 110 [arXiv:gr-qc/0512153]
- [26] E.T. Akhmedov, V. Akhmedova, D. Singleton and T. Pilling, *Int. J. Mod. Phys. A* **22** 1705 (2007)
- [27] E.T. Akhmedov, V. Akhmedova and D. Singleton, *Phys. Lett. B* **642** 124 (2006)
- [28] B. D. Chowdhury, *Pramana* **70** 3 (2008)
- [29] E.T. Akhmedov, T. Pilling and D. Singleton, *Int.J.Mod.Phys.D* **17** 2453, (2008) [arXiv:0805.2653]
- [30] V. Akhmedova, T. Pilling, A. de Gill and D. Singleton, *Phys. Lett. B* **666** 269 (2008)
- [31] V. Akhmedova, T. Pilling, A. de Gill and D. Singleton, *Phys. Lett. B* **673** 227 (2009)
- [32] K. Srinivasan and T. Padmanabhan, *Phys. Rev. D* **60** 24007 (1999)
- [33] S. Shankaranarayanan, T. Padmanabhan, and K. Srinivasan, *Class. Quant. Grav.* **19** 2671 (2002)
- [34] M. Angheben, M. Nadalini, L. Vanzo and S. Zerbini, *J.High.Energy.Phys.* **0505** (2005) 014.
- [35] D. Y. Chen and S. Z. Yang, *IJTP* **46**,2923Y (2007)
- [36] D. Y. Chen and S. Z. Yang, *New Journal of Physics* **9** 252 (2007)
- [37] R. Kerner and R. B. Mann, *Class.Quant.Grav.* **25** 095014 (2008) [arXiv:0710.0612]
- [38] R. Kerner and R. B. Mann, *Phys.Lett.B* **665** 277 (2008) [arXiv:hep-th/0803.2246]
- [39] R. Li, J. R. Ren and S. W. Wei, *Class.Quant.Grav.* **25** 125016 (2008) [arXiv:0803.1410]
- [40] R. Li and J. R. Ren, *Phys.Lett.B* **661** 370 (2008) [arXiv:0802.3954]
- [41] D. Y. Chen, Q. Q. Jiang and X. T. Zu, *Class.Quant.Grav.* **25** 205022 (2008) [arXiv:0803.3248]
- [42] D. Y. Chen, Q. Q. Jiang and X. T. Zu, *Phys.Lett.B* **665** 106 (2008) [arXiv:0804.0131]
- [43] X. X. Zeng and S. Z. Yang, *Gen.Relativ.Gravit.* **40** 2107 (2008)
- [44] K. Lin and S. Z. Yang *Acta.Phys.Sin.* **58** 744 (2009) (in Chinese)
- [45] R. D. Criscienzo and L. Vanzo, *Europhys.Lett.* **82** (2008) 60001
- [46] L. H. Li, S. Z. Yang, T. J. Zhou and R. Lin, *Europhys.Lett.* **84** (2008) 20003
- [47] Q. Q. Jiang, *Phys.Lett.B* **666** 517 (2008)
- [48] Q. Q. Jiang, *Phys.Rev.D* **78** 044009 (2008)
- [49] K. Lin and S. Z. Yang *Phys.Rev.D* **79** 064035 (2009)
- [50] K. Lin and S. Z. Yang *Phys.Lett.B* **674** 127 (2009)
- [51] F. R. Tangherlini, *Nuovo Cim.* **27** 636 (1963).
- [52] S. J. Gao and J. P. S. Lemos, *Int.J.Mod.Phys.A* **23** 2943 (2008) [arXiv:0804.0295]
- [53] R.A. Konoplya and A. Zhidenko, *Nucl.Phys.B* **777** 182, (2007)[arXiv:hep-th/0703231]
- [54] R.A. Konoplya and A. Zhidenko, *Phys.Rev.Lett.* **103** 161101 (2009) [arXiv:0809.2822]
- [55] R.A. Konoplya and A. Zhidenko, *Phys.Rev.D* **78** 104017, (2008) [arXiv:0809.2048]
- [56] R. Banerjee and B. R. Majhi, *J.High.Energy.Phys.* **0806** 095 (2008) [arXiv:0805.2220]
- [57] R. Banerjee and B. R. Majhi, *Phys.Lett.B* **674** 218 (2009) [arXiv:0808.3688]
- [58] B. R. Majhi, *Phys.Rev.D* **79** 044005 (2009) [arXiv:0809.1508]
- [59] R. Banerjee and B. R. Majhi, *Phys.Rev.D* **79** 064024 (2009) [arXiv:0812.0497]
- [60] R. Banerjee, B. R. Majhi, and D. ROY, [arXiv:0901.0466]
- [61] B. R. Majhi and S. Samanta, [arXiv:0901.2258]
- [62] R. Banerjee and B. R. Majhi, *Phys.Lett.B* **675** 243 (2009) [arXiv:0903.0250]
- [63] R. Banerjee and B. R. Majhi, *Phys.Lett.B* **662** 62 (2008) [arXiv:0801.0200]
- [64] R. Banerjee, B. R. Majhi and S. Samanta, *Phys.Rev.D* **77** 124035 (2008) [arXiv:0801.3583]
- [65] S.K. Modak, *Phys.Lett.B* **671** 167 (2009) [arXiv:0807.0959]